

KIPSONLINEEARLYPREPSESSION

BY SAEED MDCAT

www.saeedmdcat.com

FB/SAEED MDCAT

03418729745



SAEED MDCAT
SAEED MDCAT TEAM
f SAEEDMDCAT


SPECIAL THANKS TO DR.ASMA
REGARD.HUZAIFA SAEED,USAMA SOHAIL



SAEED MDCAT
SAEED MDCAT TEAM
f SAEEDMDCAT

SAEED MDCAT TEAM

f SAEEDMDCAT



SAEED MDCAT

If you want to get all the test and Lec of all
Academy then follow us on different
platforms

WEBSITE.www.saeedmdcat.com

FB Group And Page.SAEED MDCAT

INSTAGRAM.SAEED MDCAT

Twitter.Smdcat

For WhatsApp Group.03418729745

Regards.Huzaiifa Saeed,Usama Sohail

SAEED MDCAT TEAM

 SAEEDMDCAT

SAEED MDCAT

TRUE HERO ALWAYS WINS

National MDCAT 2020 Results

190+ 35 Students

185+ 218 Students

180+ 677 Students



SAEED MDCAT
SAEED MDCAT TEAM
f SAEEDMDCAT

**JOIN US FOR FREE
TO SECURE YOUR FUTURE**

Designed by: Usman Arts

f SAEEDMDCAT



QUIZZES

Practice Test Unit 1



10 Questions



7 min

Topics

Displacement, Velocity

Start Quiz

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT

www.saeedmdcat.com

FB)SAEED MDCAT

03418729745

06 : 52



1/10




7 min



Hint

Q :

A person moves along a circular track and covers half of the path as shown in figure. What will be the ratio of its distance to displacement covered?



A

 $2r$

B

 $4r$

C

 $\frac{r}{2}$

D

 $\frac{\pi}{2}$

SAEED MDCAT
SAEED MDCAT TEAM



SAEEDMDCAT

www.saeedmdcat.com

FB)SAEED MDCAT

03418729745

1

2

3

4

5

6

7

06 : 45



2/10



7 min



Hint

Q :

An athlete completes one round of a circular track of radius R in 40 sec. What will be his displacement at the end of 3 min. 20 sec?

Zero

A

B

$2\pi R$

C

$2R$

D

$7\pi R$

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT

www.saeedmdcat.com

FB)SAEED MDCAT

03418729745

1

2

3

4

5

6

7

06 : 43



3/10



7 min



Hint

Q :

A man leaves his house for a cycle ride. He comes back to his house after half-an-hour after covering a distance of one km. What is his average velocity for the ride?

A

zero

B

2 km/h

C

10 km/h

D

$\frac{1}{2}$ km/h

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT

www.saeedmdcat.com

FB)SAEED MDCAT

03418729745

1

2

3

4

5

6

7

06 : 41



4/10



7 min



Hint

Q :

Stopping distance of a moving vehicle is directly proportional to

A

square of the initial velocity

B

square of the initial acceleration

C

the initial velocity

D

the initial acceleration

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT

www.saeedmdcat.com

FB)SAEED MDCAT

03418729745

1

2

3

4

5

6

7

06 : 39



5/10



7 min



Hint

Q :

The ratio of displacement along diameter and total distance along circle:



1 : π



2 : π



π : 1



π : 2

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT

www.saeedmdcat.com

FB)SAEED MDCAT

03418729745

1

2

3

4

5

6


7

06 : 35



Q :

In 1.0 second a particle goes from point A to point B moving in a semi-circle a radius 1.0 m (figure). The magnitude of the average velocity is



3-14 m/s

A

2.0 m/s

B

1/0 m/s

C

www.saeedmdcat.com

FB) SAEED MDCAT

03418729745

1

2

3

4

5

6

7



7/10




7 min



Hint

Q :

A helicopter is flying at 3564 m above ground. If an angle of 30° is subtended at a ground point by the helicopter position 100s apart, what is the speed of the helicopter



A

100 ms⁻¹

B

150 ms⁻¹

C

20 ms⁻¹

D

25 ms⁻¹



8/10



7 min



Hint

Q :

A car travels equal distance in the same direction with velocities 60 km h^{-1} , 20 km h^{-1} and 10 km h^{-1} respectively. The average velocity of the car over the whole journey of motion is



8 ms^{-1}



6 ms^{-1}



7 ms^{-1}



5 ms^{-1}

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



9/10



7 min



Hint

Q :

Two bodies are moving in opposite directions with velocity v . The relative velocity of one with respect to the other is

A

$$\sqrt{2}v$$

B

$$2v$$

C

$$v$$

D

Zero

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



9/10



7 min



Hint

Q :

Two bodies are moving in opposite directions with velocity v . The relative velocity of one with respect to the other is

A

$$\sqrt{2}v$$

B

$$2v$$

C

$$v$$

D

Zero

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



10/10



7 min



Hint

Q :

A moves with 65 km/h while B is coming back of A with 80 km/h. The relative velocity of B with respect to A is

80 km/h

A

60 km/h

B

15 km/h

C

145 km/h

D

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



QUIZ RESULT

Practice Test Unit 1



10



7 min



18-Feb-2021



0 sec



0/10



0.0%


SAEED MDCAT

Result Detail

SAEED MDCAT TEAM




SAEEDMDCAT





Practice Test Unit 1

A person moves along a circular track and covers half of the path as shown in figure. What will be the ratio of its distance to displacement covered?



A

$2r$

B

$4r$

C

$\frac{r}{2}$

D

$\frac{\pi}{2}$

SAEED MDCAT

SAEED MDCAT TEAM

Explanation



SAEEDMDCAT

$$\frac{S}{|d|} = \frac{r\theta}{2r} = \frac{\pi}{2}$$



Practice Test Unit 1

Q :

An athlete completes one round of a circular track of radius R in 40 sec. What will be his displacement at the end of 3 min. 20 sec?

A

Zero

B

$2\pi R$

C

$2R$

D

$7\pi R$

Explanation

SAEED MDCAT TEAM

Total time of motion is 3 min 20 sec = 200 sec. As time period of circular motion is 40 sec so in 200 sec athlete will complete 5 revolution i.e., he will be at starting point i.e., displacement = zero.

1

2

3

4

5

6

7



Practice Test Unit 1



Correct



Unattempted



Incorrect



3/10

Q :

A man leaves his house for a cycle ride. He comes back to his house after half-an-hour after covering a distance of one km. What is his average velocity for the ride?



zero



2 km/h



10 km/h



$\frac{1}{2}$ km/h



Explanation

SAEEDMDCAT

Since displacement is zero.

1

2

3

4

5

6

7



Practice Test Unit 1

Q :

Stopping distance of a moving vehicle is directly proportional to

A

square of the initial velocity

B

square of the initial acceleration

C

the initial velocity

D

the initial acceleration

Explanation

Let s be the distance travelled by the vehicle before it stops. Final velocity

$$v=0$$

, initial velocity = u Using equation of motion

$$v^2 - u^2 = 2aS$$

$$0^2 - u^2 = 2aS$$

Stopping distance,

$$S = -u^2/2a$$

1

2

3

4

5

6

7



Practice Test Unit 1

Q :

The ratio of displacement along diameter and total distance along circle:

A

$$1 : \pi$$

B

$$2 : \pi$$

C

$$\pi : 1$$

D

$$\pi : 2$$

Explanation

$$|\vec{d}| = 2r \text{ (along diameter)}$$

$$S = 2\pi r \text{ (distance along circle)}$$

$$\frac{|\vec{d}|}{S} = \frac{2r}{2\pi r} = 1 : \pi$$

1

2

3

4


5

6

7



Practice Test Unit 1



A

3-14 m/s

B

2.0 m/s

C

1/0 m/s

D

zero

Explanation

$$d = 2r = 2(1) = 2$$
$$v_{ave} = \frac{\Delta d}{\Delta t} = \frac{2}{1} = 2ms^{-1}$$

1

2

3

4

5

6

7



Practice Test Unit 1



A

100 ms⁻¹

B

150 ms⁻¹

C

20 ms⁻¹

D

25 ms⁻¹

Explanation

Here $\tan 30^\circ = x/3564$

$$x = 3564 \times \frac{1}{\sqrt{3}} = 2000 \text{ m}$$

$$\text{Speed of the helicopter} = \frac{2000}{100} = 20 \text{ ms}^{-1}$$

4

5

6

7

8

9

10



Practice Test Unit 1

Q :

A car travels equal distance in the same direction with velocities 60 km h^{-1} , 20 km h^{-1} and 10 km h^{-1} respectively. The average velocity of the car over the whole journey of motion is

A

8 ms^{-1}

B

6 ms^{-1}

C

7 ms^{-1}

D

5 ms^{-1}

Explanation

Average velocity

$$= \frac{\frac{3x}{60} + \frac{3x}{20} + \frac{3x}{10}}{\frac{x}{60} + \frac{x}{20} + \frac{x}{10}} = \frac{\frac{3x}{60}}{\frac{x+3x+6x}{60}} = \frac{3x \times 60}{10x} = 18$$

4

5

6

7

8

9

10



Practice Test Unit 1

Q :

A car travels equal distance in the same direction with velocities 60 km h^{-1} , 20 km h^{-1} and 10 km h^{-1} respectively. The average velocity of the car over the whole journey of motion is

A

8 ms^{-1}

B

6 ms^{-1}

C

7 ms^{-1}

D

5 ms^{-1}

Explanation



SAEEDMDCAT

$$= \frac{3x}{\frac{x+3x+6x}{60}} = \frac{3x \times 60}{10x} = 18 \text{ km h}^{-1} = 5 \text{ ms}^{-1}$$

4

5

6

7

8

9

10



Practice Test Unit 1

Two bodies are moving in opposite directions with velocity v . The relative velocity of one with respect to the other is

A

$$\sqrt{2}v$$

B

$$2v$$

C

$$v$$

D

Zero

Explanation

$$v_{rel} = \sqrt{v_1^2 + v_2^2 - 2v_1v_2\cos\theta}$$

$$v_{rel} = \sqrt{v^2 + v^2 - 2v^2\cos 180^\circ}$$

$$V_{rel} = \sqrt{v^2 + v^2 - 2v^2} = \sqrt{4v^2} = 2v$$

4

5

6

7

8

9

10



Practice Test Unit 1



Incorrect



10/10

Q :

A moves with 65 km/h while B is coming back of A with 80 km/h. The relative velocity of B with respect to A is

80 km/h

A

60 km/h

B

15 km/h

C

145 km/h

D

SAEED MDCAT
SAEED MDCAT TEAM



Explanation

SAEEDMDCAT

$$v_B + v_A = v_B + v_A = 80 + 65 = 145 \text{ km/hr}$$

4

5

6

7

8

9

10

www.saeedmdcat.com



FB) SAEED MDCAT

03418729745



QUIZZES

Practice test 2 unit 1



10 Questions



10 min

Topics

Acceleration, Graphical representation of
acceleration with velocity time graph

Start Quiz

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



1/10



10 min



Hint

Q :

A constant force F changes the velocity of a 80kg sprinter from 3ms^{-1} to 4ms^{-1} in 0.5 sec. The acceleration of sprinter will be

A

1.5ms^{-2}

B

2.5ms^{-2}

C

2.0ms^{-2}

D

3.0ms^{-2}

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



2/10




10 min



Hint

Q : A ball is projected upwards with an initial velocity $V_i = 100 \text{ ms}^{-1}$ as shown in the figure. It comes back after sometime and strikes ground with the same velocity but pointing downward. What is the angle between acceleration when moving upward to acceleration coming down?



A

180°

B

90°

C

60°

D

0°

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



3/10



10 min



Hint

Q :

A baseball is thrown vertically into the air. The acceleration of the ball at its highest point is:



Zero



g , up



g , down



$2g$, down

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



4/10



10 min



Hint

Q :

A passenger in a moving train tosses a coin. If the coin falls behind him, the train must be moving with



A An acceleration



B a deceleration



C A uniform speed



D any of the above

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



5/10




10 min



Hint

Q :

The diagram shows a velocity-time graph for a particle moving along a straight line. What is the displacement of the particle between 0 s and 25 s?



A

90 m

B

130 m

C

120

D

150 m



SAEEDMDCAT



6/10




10 min



Hint

Q : The graph shows how the speed of an object changes with time. Which section of the graph shows the object moving with an increasing acceleration?



A



B



C



D



SAEEDMDCAT



7/10




10 min




Hint

Q : Acceleration-time graph of a body is shown.
The Corresponding velocity time graph of the
same body is



A



B




SAEEDMDCAT




03418729745

A



B



C



D





8/10



10 min



Hint

Q : Displacement time graph of a ball thrown vertically upward is shown in figure then its v-t graph is:


Displacement



A



B



C



D





9/10




10 min



Hint

Q : In the given graph distance traveled is:



A

200m

B

300 m

C

250 m

D

400m

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



10/10




10 min




Hint

Q : A batsman hits a sixer and the ball reaches out of the cricket ground. Which of the following graphs describes the variation of the cricket ball's vertical velocity v with time between t_1 (the time of hitting the bat and time t_2 (the time of touching the ground)?

A



B




C




t_2 (the time of touching the ground)?

03418729745


A



B



C



D





QUIZ RESULT

Practice test 2 unit 1



10



10 min



19-Feb-2021



0 sec



0/10



0.0%


SAEED MDCAT

Result Detail

SAEED MDCAT TEAM



SAEEDMDCAT





Practice test 2 unit 1



Correct



Unattempted



Incorrect



1/10

Q:

A constant force F changes the velocity of a 80kg sprinter from 3ms^{-1} to 4ms^{-1} in 0.5 sec. The acceleration of sprinter will be

A

1.5ms^{-2}

B

2.5ms^{-2}

C

2.0ms^{-2}

D

3.0ms^{-2}

Explanation



SAEEDMDCAT

$$a = \frac{v_f - v_i}{t} = \frac{4 - 3}{0.5} = 2\text{ms}^{-2}$$

1

2

3

4

5


6

7



Practice test 2 unit 1

Q : A ball is projected upwards with an initial velocity $V_i = 100 \text{ ms}^{-1}$ as shown in the figure. It comes back after sometime and strikes ground with the same velocity but pointing downward. What is the angle between acceleration when moving upward to acceleration coming down?



A 180°

B 90°

C 60°

D 0°

SAEED MDCAT

SAEED MDCAT TEAM

Explanation



SAEEDMDCAT

Direction of acceleration due to gravity toward earth center.



Practice test 2 unit 1



Correct



Unattempted



Incorrect



3/10

Q :

A baseball is thrown vertically into the air. The acceleration of the ball at its highest point is:

A

Zero

B

g , up

C

g , down

D

$2g$, down

Explanation

The acceleration of ball at highest point when thrown vertically upward option (c) g , down.

1

2

3

4

5

6

7



Practice test 2 unit 1



Correct



Unattempted



Incorrect



4/10

Q:

A passenger in a moving train tosses a coin. If the coin falls behind him, the train must be moving with



An acceleration



a deceleration



A uniform speed



any of the above

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT

1

2

3

4

5


6

7



Practice test 2 unit 1

the displacement of the particle between 0 s and 25 s?



A 90 m

B 130 m

C 120

D 150 m

Explanation

Explanation Displacement = area under graph

$$= (2 \times 5) + \frac{1}{2} (25 - 5)(2 + 10)$$

$$= 130 \text{ m}$$



Practice test 2 unit 1



Correct



Unattempted



Incorrect



6/10

Q : The graph shows how the speed of an object changes with time. Which section of the graph shows the object moving with an increasing acceleration?



A

A

B

B

C

C

D

D

1

2

3

4

5

6

7



Practice test 2 unit 1



Correct



Unattempted




Incorrect




7/10

Q : Acceleration-time graph of a body is shown.
The Corresponding velocity time graph of the
same body is



A



B



SAEEDMDCAT

1

2

3

4

5


6

7




Practice test 2 unit 1

C



D



Explanation

Rate of change of velocity is acceleration.

We know that $a = \frac{\Delta v}{\Delta t}$ where a is acceleration and v is velocity. In the given graph acceleration is constant for first part of motion. So, velocity increases uniformly.

When $a=0$ i.e, $0 = \frac{\Delta v}{\Delta t} \Rightarrow v = \text{constant}$ Thus, we can draw velocity time graph as:






Practice test 2 unit 1


A




B



C



D



SAEED MDCAT

Explanation

SAEED MDCAT TEAM

Explanation It v-t graph is:



4

5

6

7

8

9

10



Practice test 2 unit 1



Correct



Unattempted




Incorrect



9/10

Q : In the given graph distance traveled is:



200m



300 m



250 m



400m

Explanation

S=Area under the graph, S= Area of

$$S = \frac{1}{2} (30 + 10)(10) = 200m$$

trapezium,

4

5

6

7

8

9

10



Practice test 2 unit 1



Correct



Unattempted




Incorrect



10/10

Q : A batsman hits a sixer and the ball reaches out of the cricket ground. Which of the following graphs describes the variation of the cricket ball's vertical velocity v with time between t_1 (the time of hitting the bat and time t_2 (the time of touching the ground)?

A



B



4

5

6

7

8


9

10




Practice test 2 unit 1


B



C



D



Explanation

Initially the ball has vertical component of motion upward. It decreases, becomes zero and then begins to increase in downward direction.

www.saeedmdcat.com



FB) SAEED MDCAT

03418729745



QUIZZES

Practice test 3 unit 1



10 Questions



7 min

Topics

Newton's laws of motion, Linear Momentum, Law of conservation of momentum

Start Quiz

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



1/10



7 min



Hint

Q :

The engine of a car produces an acceleration of 6 ms^{-2} in the car. If this car pulls another car of the same mass, then the acceleration would be

A

6 ms^{-2}

B

12 ms^{-2}

C

3 ms^{-2}

D

1.5 ms^{-2}

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



2/10



7 min



Hint

Q :

What is the resultant force shown in fig?



A

7 N towards

B

17 N in arbitrary direction

C

3 N toward left

D

10 N towards left

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



3/10




7 min



Hint

Q : Two masses of 4 kg and 5 kg are connected by a string passing through a frictionless pulley and are kept on a frictionless table as shown in the figure.



The acceleration of 5 kg mass is

A

49m/s²

B

5.44m/s²

C

19.5m/s²

D

2.72m/s²



4/10



7 min



Hint

Q : A particle of mass "m" is projected with a velocity $6\hat{i} + 8\hat{j}$. Find the change in momentum when it touches ground:

A

16 m

B

12 m

C

10 m

D

20 m

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



5/10



7 min



Hint

Q :

Two masses of 1 gm and 4 gm are moving with the same kinetic energy. The ratio of their linear momentum will be



1:16



1:2



$\sqrt{2}:1$



4:1

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



6/10



7 min



Hint

Q : When the velocity is doubled:

A

K.E is doubled

B

Acceleration is doubled

C

Momentum is doubled

D

P.E is doubled

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT

1

2

3

4

5

6

7



7/10



7 min



Hint

Q :

The momentum is most closely related to

A

Force

B

Impulse

C

Power

D

K.E.

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



8/10



7 min



Hint

Q :

A wagon weighing 1000 kg is moving with a velocity 50 km/h on smooth horizontal rails. A mass of 250 kg is dropped into it. The velocity with which it moves now is

A

40 km/hour

B

50 km/hour

C

2.5 km/hour

D

20 km/hour

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



9/10



7 min



Hint

Q : In which of the following cases forces may not be required to keep the

A

Particle going in a circle

B

Particle going along a straight line

C

The momentum of the particle constant

D

Acceleration of the particle constant

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



10/10



7 min



Hint

Q : The motion of a rocket is based on the principle of conservation of

A

Mass

B

Kinetic energy

C

Linear momentum

D

Angular momentum

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



QUIZ RESULT

Practice test 3 unit 1



10



7 min



19-Feb-2021



0 sec



0/10



0.0%


SAEED MDCAT

Result Detail

SAEED MDCAT TEAM



SAEEDMDCAT





Practice test 3 unit 1

Q :

The engine of a car produces an acceleration of 6 ms^{-2} in the car. If this car pulls another car of the same mass, then the acceleration would be

A

6 ms^{-2}

B

12 ms^{-2}

C

3 ms^{-2}

D

1.5 ms^{-2}

Explanation

Force applied by engine = 6 ms^{-2}

When two cars are pulled,

$$(m + m)a = 6 \text{ ms}^{-2}$$

$$2ma = 6 \text{ ms}^{-2} \text{ or } a = 3 \text{ ms}^{-2}$$



Practice test 3 unit 1



Correct



Unattempted




Incorrect



2/10

Q:

What is the resultant force shown in fig?



A

7 N towards

B

17 N in arbitrary direction

C

3 N toward left

D

10 N towards left

Explanation



SAEEDMDCAT

According to 2nd law of motion.

$$F_{net} = 10 + 2 - 5 = 3 \text{ N towards left}$$

1

2

3

4

5


6

7



Practice test 3 unit 1

by a string passing through a frictionless pulley and are kept on a frictionless table as shown in the figure.



The acceleration of 5 kg mass is

A

49m/s²

B

5.44m/s²

C

19.5m/s²

D

2.72m/s²

SAEED MDCAT

SAEED MDCAT TEAM

Explanation



SAEEDMDCAT

$$a = \frac{m_2}{m_1 + m_2} \times g = \frac{5}{4 + 5} \times 9.8 = \frac{49}{9} = 5.44 \text{ ms}$$



Practice test 3 unit 1



Correct



Unattempted



Incorrect



4/10

Q : A particle of mass "m" is projected with a velocity $6\hat{i} + 8\hat{j}$. Find the change in momentum when it touches ground:



16 m



12 m



10 m



20 m

Explanation

$$P = mv \quad |\vec{v}| = \sqrt{6^2 + 8^2} = 10$$
$$P = 10m$$

1

2

3

4

5

6

7



Practice test 3 unit 1



Correct



Unattempted



Incorrect



5/10

Q :

Two masses of 1 gm and 4 gm are moving with the same kinetic energy. The ratio of their linear momentum will be



1:16



1:2



$\sqrt{2}:1$



4:1

SAEED MDCAT

SAEED MDCAT TEAM

Explanation



SAEEDMDCAT

$$P = \sqrt{2m K.E} \Rightarrow \frac{P_1}{P_2} = \sqrt{\frac{1}{4} \times \frac{E}{E}} = \frac{1}{2}$$

1

2

3

4

5

6

7



Practice test 3 unit 1



Correct



Unattempted



Incorrect



6/10

Q : When the velocity is doubled:



K.E is doubled



Acceleration is doubled



Momentum is doubled



P.E is doubled

Explanation

$$P = mv, P' = m(2v) = 2P$$



SAEEDMDCAT

1

2

3

4

5

6

7



Practice test 3 unit 1



Correct



Unattempted



Incorrect



7/10

Q:

The momentum is most closely related to



Force



Impulse



Power



K.E.

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT

4

5

6

7

8

9

10



Practice test 3 unit 1

A wagon weighing 1000 kg is moving with a velocity 50 km/h on smooth horizontal rails. A mass of 250 kg is dropped into it. The velocity with which it moves now is

A 40 km/hour

B 50 km/hour

C 2.5 km/hour

D 20 km/hour

Explanation

$$m_1 = 1000 \text{ kg}$$

$$v_1 = 50 \text{ kmph}$$

$$m_2 = 1000 + 250 = 1250 \text{ kg.}$$

$$v_2 = ?$$

by principle of conservation of momentum,

$$m_1 v_1 = m_2 v_2$$

$$v_2 = \{(m_1 v_1) / (m_2)\} = \{(1000 \times 50) / (1250)\}$$

$$v_2 = 40 \text{ kmph}$$



Practice test 3 unit 1



Correct



Unattempted



Incorrect



9/10

Q : In which of the following cases forces may not be required to keep the

A

Particle going in a circle

B

Particle going along a straight line

C

The momentum of the particle constant

D

Acceleration of the particle constant

Explanation

If momentum remains constant then force will be zero because

$$F = \frac{\Delta p}{\Delta t}$$

4

5

6

7

8

9

10



Practice test 3 unit 1



Correct



Unattempted



Incorrect



10/10

Q : The motion of a rocket is based on the principle of conservation of



Mass



Kinetic energy



Linear momentum



Angular momentum

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT

4

5

6

7

8

9

10

www.saeedmdcat.com



FB) SAEED MDCAT

03418729745



QUIZZES

Practice test 4 unit 1



10 Questions



7 min

Topics

Collision, Elastic collision, Elastic collision in one dimension under different cases

Start Quiz

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



1/10



7 min



Hint

Q :

A billiard ball moving with a speed of 5 m/s collide with an identical ball, originally at rest. If first ball stops dead after the collision, then second ball will move forward with speed of

A

10 m/s

B

5 m/s

C

2.5 m/s

D

9 m/s

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



2/10




7 min



Hint

Q : When a very heavy ball 'B1' collide with a stationary target 'B2' of negligible mass, after collision the final velocity of ball 'B2' will



A

become zero

B

become half

C

become doubled as compared to B1

D

same as the B1

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



3/10



7 min



Hint

Q : A 10 kg object collides with stationary 5 kg object and after collision they stick together and move forward with velocity 4ms^{-1} . What is the velocity with which the 10 kg object hit the second one

A

6ms^{-1}

B

8ms^{-1}

C

12ms^{-1}

D

10ms^{-1}

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



4/10



7 min



Hint

Q :

When a ball bounces back from floor such that sound and heat is produced then collision

A

Must be elastic

B

Must be inelastic

C

Momentum is conserved but K.E is not conserved

D

Both B and C

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT

1

2

3

4

5

6

7



5/10



7 min



Hint

Q :

A ball of mass 2 kg travelling at 8 ms^{-1} strikes a ball of mass 4 kg travelling at 2 ms^{-1} . Both balls are moving along the same straight line as shown



After collision, both balls move at the same velocity v . What is the magnitude of the velocity v ?

A

4 ms^{-1}

B

5 ms^{-1}

C

6 ms^{-1}

D

8 ms^{-1}



6/10




7 min



Hint

Q :

Two similar spheres, each of mass m and travelling with speed v , are moving towards each other.



The spheres have a head on elastic collision. Which statement is correct?

The spheres stick together on impact

A

The total kinetic energy after impact is mv^2

B

The total kinetic energy before impact is zero

C

1

2

3


4

5

6

7

Two similar spheres, each of mass m and travelling with speed v , are moving towards each other.



The spheres have a head on elastic collision. Which statement is correct?

A

The spheres stick together on impact

B

The total kinetic energy after impact is mv^2

C

The total kinetic energy before impact is zero

D

The total momentum before impact is $2mv$



7/10



7 min



Hint

Q :

Two railway trucks of masses m and $3m$ move towards each other in opposite directions with speeds $2v$ and v respectively. These trucks collide and stick together.

What is the speed of the trucks after the collision?

A

$$\frac{v}{4}$$

B

$$\frac{v}{2}$$

C

$$v$$

D

$$\frac{5v}{4}$$

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT

1

2

3

4

5

6

7

03418729745

Q :

Which one of the following is true in the case of inelastic collision?

A

Total energy
Energy

Conserved

Momentum

Conserved

Kinetic

Conserved

B

Total energy
Energy

Conserved
conserved

Momentum

Conserved

Kinetic

Not

C

Total energy
Energy

Conserved
Conserved

Momentum

Not conserved

Kinetic

D

Total energy
Energy

Not conserved
conserved

Momentum

Conserved

Kinetic

Not



9/10



7 min



Hint

Q :

A ball is dropped from a height of 10 m. It is embedded 1 m in sand and stops. In this process

A

only momentum is conserved

B

only kinetic energy is conserved

C

both momentum and kinetic energy are conserved

D

neither momentum nor kinetic energy is conserved

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



10/10



7 min



Hint

Q :

When two bodies collide elastically then the quantity conserved is



Kinetic energy



Momentum



Both



None

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



QUIZ RESULT

Practice test 4 unit 1



10



7 min



19-Feb-2021



0 sec



0/10



0.0%


SAEED MDCAT

Result Detail

SAEED MDCAT TEAM



SAEEDMDCAT





Practice test 4 unit 1



Correct



Unattempted



Incorrect



1/10

Q :

A billiard ball moving with a speed of 5 m/s collide with an identical ball, originally at rest. If first ball stops dead after the collision, then second ball will move forward with speed of

A

10 m/s

B

5 m/s

C

2.5 m/s

D

9 m/s

SAEED MDCAT

SAEED MDCAT TEAM

Explanation



SAEEDMDCAT

When two identical balls collide elastically their velocities are interchange.

1

2

3

4

5

6

7



Practice test 4 unit 1




Incorrect



2/10

Q : When a very heavy ball 'B1' collide with a stationary target 'B2' of negligible mass, after collision the final velocity of ball 'B2' will



A

become zero

B

become half

C

become doubled as compared to B1

D

same as the B1

Explanation



SAEEDMDCAT

$$m_1 \gg m_2, v_2 = 0$$

$$v_1' = v_1, v_2' = 2v_1$$

1

2

3

4

5

6

7



Practice test 4 unit 1



Incorrect



3/10

Q : A 10 kg object collides with stationary 5 kg object and after collision they stick together and move forward with velocity 4ms^{-1} . What is the velocity with which the 10 kg object hit the second one

A

6ms^{-1}

B

8ms^{-1}

C

12ms^{-1}

D

10ms^{-1}

Explanation

$$m_1 v_1 = (m_1 + m_2) v_c$$

$$v_1 = \left(\frac{m_1 + m_2}{m_1} \right) v_c$$

$$= \frac{15}{10} \times 4 = 6\text{ms}^{-1}$$

1

2

3

4

5

6

7



Practice test 4 unit 1



Correct



Unattempted



Incorrect



4/10

Q :

When a ball bounces back from floor such that sound and heat is produced then collision

A

Must be elastic

B

Must be inelastic

C

Momentum is conserved but K.E is not conserved

D

Both B and C



Explanation

SAEEDMDCAT

Application of elastic collision

1

2

3

4

5

6

7



Practice test 4 unit 1

Q :

A ball of mass 2 kg travelling at 8 ms^{-1} strikes a ball of mass 4 kg travelling at 2 ms^{-1} . Both balls are moving along the same straight line as shown



After collision, both balls move at the same velocity v . What is the magnitude of the velocity v ?

A

4 ms^{-1}

B

5 ms^{-1}

C

6 ms^{-1}

D

8 ms^{-1}



SAEEDMDCAT

Explanation



Practice test 4 unit 1

The spheres have a head on elastic collision.
Which statement is correct?

A

The spheres stick together on impact

B

The total kinetic energy after impact is mv^2

C

The total kinetic energy before impact is zero

D

The total momentum before impact is $2mv$

Explanation



SAEEDMDCAT

Kinetic energy is conserved for elastic collision.

1

2

3

4

5

6

7



Practice test 4 unit 1

What is the speed of the trucks after the collision?

A

$$\frac{v}{4}$$

B

$$\frac{v}{2}$$

C

$$v$$

D

$$\frac{5v}{4}$$

Explanation

This is a perfectly inelastic collision.



$$m(2v) + 3m(-v) = (m + 3m)v_f$$

$$-mv = 4mv_f$$

$$v_f = -\frac{1}{4}v$$



Practice test 4 unit 1

What is the speed of the trucks after the collision?

A

$$\frac{v}{4}$$

B

$$\frac{v}{2}$$

C

$$v$$

D

$$\frac{5v}{4}$$

Explanation

perfectly inelastic collision.



$$-mv = (m + 3m)v_t$$

1



SAEEDMDCAT



Practice test 4 unit 1

A

Total energy
Energy

Momentum

Kinetic

Conserved

Conserved

Conserved

B

Total energy
Energy

Momentum

Kinetic

Conserved
conserved

Conserved

Not

C

Total energy
Energy

Momentum

Kinetic

Conserved
Conserved

Not conserved

D

Total energy
Energy

Momentum

Kinetic

Not conserved
conserved

Conserved

Not

SAEED MDCAT

SAEED MDCAT TEAM

Explanation



SAEEDMDCAT

Total momentum and total energy remain conserved in all types of collision.

4

5

6

7

8

9

10



Practice test 4 unit 1



Incorrect



9/10

Q :

A ball is dropped from a height of 10 m. It is embedded 1 m in sand and stops. In this process

A

only momentum is conserved

B

only kinetic energy is conserved

C

both momentum and kinetic energy are conserved

D

neither momentum nor kinetic energy is conserved

SAEED MDCAT TEAM

Explanation



SAEEDMDCAT

During inelastic collision only momentum is conserved

4

5

6

7

8

9

10



Practice test 4 unit 1



Incorrect



10/10

Q :

When two bodies collide elastically then the quantity conserved is



Kinetic energy



Momentum



Both



None

SAEED MDCAT

SAEED MDCAT TEAM

Explanation



SAEEDMDCAT

When bodies collide elastically both momentum and K.E remain conserved.

4

5

6

7

8

9

10

www.saeedmdcat.com



FB) SAEED MDCAT

03418729745



QUIZZES

Practice test 5 unit # 1



10 Questions



7 min

Topics

Projectile motion

Start Quiz

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



1/10



7 min



Hint

Q :

A cricket ball is hit at 45° to the horizontal with a kinetic energy K . The kinetic energy at the highest point is

A

0

B

$K/2$

C

$K/\sqrt{2}$

D

K

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



2/10



7 min



Hint

Q :

A bomber drops a bomb, when it is vertically above the target. Its misses the target because



Vertically component of the velocity of the bomber



Force of gravity



Acceleration of the bomber



Horizontal component of the velocity of the bomber

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



3/10




7 min



Hint

Q :

A parabolic path for a projectile is shown in the figure. At which point acceleration is minimum?



B



A



C



Not possible

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



4/10



7 min



Hint

Q :

A body is thrown horizontally from the top of a tower of height 5 m. It touches the ground at a distance of 10 m from the foot of the tower. The initial velocity of the body is ($g = 10 \text{ ms}^{-2}$)

A

2.5 ms^{-1}

B

5 ms^{-1}

C

10 ms^{-1}

D

20 ms^{-1}

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



5/10



7 min



Hint

Q :

A ball is thrown horizontally from the top of a tower. What happens to the horizontal component of its acceleration?

A

First increases and then decreases

B

Increase

C

Decrease

D

Zero



SAEEDMDCAT



6/10



7 min



Hint

Q :

At maximum height in projectile motion the horizontal component of velocity will



Maximum



Zero



Constant



Both (b) and (c)

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



7/10



7 min



Hint

Q : An aero plane is flying horizontally with a velocity of 10 m/s and at a height of 1960 m . When it is vertically above a point A on the ground, a bomb is released from it. The bomb strikes the ground at point B. The distance AB is (ignoring air resistance)

A

100 m

B

200 m

C

400 m

D

2 km

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



8/10



7 min



Hint

Q :

A projectile is fired horizontally from a 490m high cliff with a velocity of 80ms^{-1} . The time taken by the projectile to reach the ground is

A

10 sec

B

7.5 sec

C

2.5 sec

D

5 sec

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



9/10



7 min



Hint

Q : A man projects a coin upwards from the gate of a uniformly moving train. The path of coin for the man will be

A

Parabolic

B

Inclined straight line

C

Vertical straight line

D

Horizontal straight line

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



10/10



7 min



Hint

Q :

For projectile motion in the absence of air resistance:



Vertical speed is constant



Horizontal force is constant



Horizontal acceleration is zero



Vertical acceleration is zero

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



QUIZ RESULT

Practice test 5 unit # 1



10



7 min



19-Feb-2021



0 sec



0/10



0.0%

SAEED MDCAT

Result Detail

SAEED MDCAT TEAM



SAEEDMDCAT



Practice test 5 unit # 1



Incorrect



1/10

Q :

A cricket ball is hit at 45° to the horizontal with a kinetic energy K . The kinetic energy at the highest point is

A

0

B

$K/2$

C

$K/\sqrt{2}$

D

K

Explanation

Kinetic energy at highest point,

$$K_H = K \cos^2 45^\circ = \frac{K}{2}$$

1

2

3

4

5

6

7



Practice test 5 unit # 1



Incorrect



2/10

Q :

A bomber drops a bomb, when it is vertically above the target. Its misses the target because



Vertically component of the velocity of the bomber



Force of gravity



Acceleration of the bomber



Horizontal component of the velocity of the bomber

SAEED MDCAT TEAM

Explanation



SAEEDMDCAT

Due to air friction, horizontal velocity of bomb changes

1

2

3

4

5

6

7



Practice test 5 unit # 1



Incorrect



3/10

Q :

A parabolic path for a projectile is shown in the figure. At which point acceleration is minimum?



B



A



C



Not possible

SAEED MDCAT TEAM

Explanation



SAEEDMDCAT

For whole projectile trajectory
 $a=g$

1

2

3

4

5

6

7



Practice test 5 unit # 1



Incorrect



4/10

Q :

A body is thrown horizontally from the top of a tower of height 5 m. It touches the ground at a distance of 10 m from the foot of the tower. The initial velocity of the body is ($g = 10 \text{ ms}^{-2}$)



2.5 ms^{-1}



5 ms^{-1}



10 ms^{-1}



20 ms^{-1}

SAEED MDCAT

SAEED MDCAT TEAM

Explanation



SAEEDMDCAT

$$S = v \times \sqrt{\frac{2h}{g}} \Rightarrow 10 = v \times \sqrt{2 \times \frac{5}{10}} \Rightarrow v = 10$$

1

2

3

4

5

6

7



Practice test 5 unit # 1

Q :

A ball is thrown horizontally from the top of a tower. What happens to the horizontal component of its acceleration?

A

First increases and then decreases

B

Increase

C

Decrease

D

Zero

Explanation



SAEEDMDCAT

For projectile motion: $F_x = 0$, $a_x = 0$

1

2

3

4

5

6

7



Practice test 5 unit # 1



Correct



Unattempted



Incorrect



6/10

Q :

At maximum height in projectile motion the horizontal component of velocity will



Maximum



Zero



Constant



Both (b) and (c)

SAEED MDCAT

SAEED MDCAT TEAM

Explanation



SAEEDMDCAT

In projectile motion the horizontal component of velocity will remain constant

1

2

3

4

5

6

7



Practice test 5 unit # 1



Incorrect



7/10

Q : An aero plane is flying horizontally with a velocity of 10 m/s and at a height of 1960 m. When it is vertically above a point A on the ground, a bomb is released from it. The bomb strikes the ground at point B. The distance AB is (ignoring air resistance)

A

100 m

B

200 m

C

400 m

D

2 km

Explanation

$$AB = R = u \sqrt{\frac{2H}{g}} = 10 \times \sqrt{\frac{2 \times 1960}{9.8}}$$

$$R = 10 \times 20 = 200 \text{ m}$$

1

2

3

4

5

6

7



Practice test 5 unit # 1



Incorrect



8/10

Q :

A projectile is fired horizontally from a 490m high cliff with a velocity of 80ms^{-1} . The time taken by the projectile to reach the ground is

A

10 sec

B

7.5 sec

C

2.5 sec

D

5 sec

Explanation

$$y = \frac{1}{2}gt^2$$

$$t = \sqrt{\frac{2y}{g}} = \sqrt{\frac{2 \times 490}{9.8}} = 10 \text{ sec}$$

4

5

6

7

8

9

10



Practice test 5 unit # 1



Correct



Unattempted



Incorrect



9/10

Q : A man projects a coin upwards from the gate of a uniformly moving train. The path of coin for the man will be

A

Parabolic

B

Inclined straight line

C

Vertical straight line

D

Horizontal straight line

Explanation

Because horizontal velocity is same for coin and the observer. So relative horizontal displacement will be zero.

4

5

6

7

8

9

10



Practice test 5 unit # 1



Correct



Unattempted



Incorrect



10/10

Q :

For projectile motion in the absence of air resistance:

A

Vertical speed is constant

B

Horizontal force is constant

C

Horizontal acceleration is zero

D

Vertical acceleration is zero

Explanation

$$F_x = ma_x = 0$$

$$a_x = 0$$

$$v_x = \text{constant}$$

4

5

6

7

8

9

10

www.saeedmdcat.com



FB) SAEED MDCAT

03418729745



QUIZZES

Practice test 6 unit # 1



10 Questions



7 min

Topics

Characteristics of projectile motion, Time of flight, Maximum height, Horizontal range, Effect of Air Resistance in Projectile Motion

Start Quiz

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



1/10



7 min



Hint

Q : A body of mass 1 kg is thrown with a velocity of 10 ms^{-1} at an angle of 60° with the horizontal. Its momentum at the highest point is



A 2 kg ms^{-1}



B 3 kg ms^{-1}



C 4 kg ms^{-1}



D 5 kg ms^{-1}

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



2/10



7 min



Hint

Q :

A cannonball is fired and follows the parabolic path shown below. Air resistance is negligible. Point B is the highest point on the path and points A and C are at the same height.



How do the speeds of the cannonball at the there points compare?

A

$$v_A < v_B < v_C$$

B

$$v_C < v_B < v_A$$

C

$$v_B < v_A < v_C$$

D

$$v_B < v_A = v_C$$



3/10



7 min



Hint

Q : A projectile is projected at angle of 30° with an initial velocity of 10 m/sec. It will go upto height of _____



10 m



2.5 m



1.25 m



0.75 m

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



4/10



7 min



Hint

Q : The angle of projection for which the maximum height and the horizontal range of a projectile are equal to

A

45°

B

$\tan^{-1}\left(\frac{1}{4}\right)$

C

$\tan^{-1}(4)$

D

None of these

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



5/10



7 min



Hint

Q : If a projectile is thrown with 19.6m/s velocity at 30° with x-axis, time taken to reach highest point?



1 sec



2 sec



3 sec



4 sec

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



6/10



7 min



Hint

Q : A missile is fired for maximum range with a initial velocity of 20 m/s. If $g=10\text{m/s}^2$, the range of the missile is



30 m



20 m



40 m



60 m

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



7/10



7 min



Hint

Q : If the initial speed of a projectile is doubled:

A

Its range will double

B

Its range will quadruple

C

Its range will be decreased by a factor of two

D

Its range will decrease by a factor of four

SAEED MDCAT
SAEED MDCAT TEAM



SAEEDMDCAT

1

2

3

4

5

6

7



8/10



7 min



Hint

Q : The path of projectile in presence of air friction is more likely to be

A

Parabolic

B

Circle

C

Elliptical

D

Straight line

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



9/10



7 min



Hint

Q : In presence of air resistance, which one will be true for horizontal component of acceleration of projectile

A

$$a_x = 0$$

B

$$a_x = g$$

C

$$a_x \neq 0$$

D

$$a_x = -F_x$$

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



10/10



7 min



Hint

Q :

If a projectile is projected with initial speed 20 msec-1 at angle 75° with horizontal in presence of air friction, then which of the following may be the range of projectile

A

36 m

B

27 m

C

20 m

D

17 m

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



QUIZ RESULT

Practice test 6 unit # 1



10



7 min



22-Feb-2021



0 sec



0/10



0.0%


SAEED MDCAT

Result Detail

SAEED MDCAT TEAM



SAEEDMDCAT





Practice test 6 unit # 1



Correct



Unattempted



Incorrect



1/10

Q : A body of mass 1 kg is thrown with a velocity of 10 ms^{-1} at an angle of 60° with the horizontal. Its momentum at the highest point is

A

2 kg ms^{-1}

B

3 kg ms^{-1}

C

4 kg ms^{-1}

D

5 kg ms^{-1}

Explanation

SAEED MDCAT TEAM

$$P = mv \cos \theta$$

$$= 1 \times 10 \times \cos 60^\circ = 10 \left(\frac{1}{2} \right) \text{ kg ms}^{-1}$$

$$= 5 \text{ kg ms}^{-1}$$

1

2

3

4

5

6

7



Practice test 6 unit # 1



Correct



Unattempted




Incorrect



2/10

Q :

A cannonball is fired and follows the parabolic path shown below. Air resistance is negligible. Point B is the highest point on the path and points A and C are at the same height.



How do the speeds of the cannonball at the there points compare?



$v_A < v_B < v_C$



$v_C < v_B < v_A$



$v_B < v_A < v_C$



$v_B < v_A = v_C$

1

2

3

4

5

6

7



Practice test 6 unit # 1



Correct



Unattempted



Incorrect



3/10

Q : A projectile is projected at angle of 30° with an initial velocity of 10 m/sec. It will go upto height of _____



10 m



2.5 m



1.25 m



0.75 m

Explanation

$$H = \frac{v_i^2 \sin^2 \theta}{2g} = \frac{100 \times \left(\frac{1}{2}\right)^2}{2 \times 10} = \frac{25}{20} = 1.25 \text{ m}$$

1

2

3

4

5

6

7



Practice test 6 unit # 1



Correct



Unattempted



Incorrect



4/10

Q : The angle of projection for which the maximum height and the horizontal range of a projectile are equal to



45o



$\tan^{-1}(\frac{1}{4})$



$\tan^{-1}(4)$



None of these

Explanation

$$H = \frac{R}{4} \tan \theta$$

$$\tan^{-1}(4) = \theta$$

1

2

3

4

5

6

7



Practice test 6 unit # 1



Correct



Unattempted



Incorrect



5/10

Q : If a projectile is thrown with 19.6m/s velocity at 30° with x-axis, time taken to reach highest point?



1 sec



2 sec



3 sec



4 sec

Explanation

As the formula to reach maximum height is:

$$t = \frac{v_i \sin \theta}{g} = \frac{19.6 \times \sin 30}{9.8} = 1s$$

1

2

3

4

5

6

7



Practice test 6 unit # 1



Correct



Unattempted



Incorrect



6/10

Q : A missile is fired for maximum range with a initial velocity of 20 m/s. If $g=10\text{m/s}^2$, the range of the missile is



30 m



20 m



40 m



60 m

Explanation

$$R_{\max} = \frac{v_i^2}{g} = \frac{20 \times 20}{10} = 40\text{m}$$

1

2

3

4

5

6

7



Practice test 6 unit # 1

Q : If the initial speed of a projectile is doubled:

A

Its range will double

B

Its range will quadruple

C

Its range will be decreased by a factor of two

D

Its range will decrease by a factor of four

Explanation

As $R \propto v_i^2$ so by doubling the initial velocity v_i Range will quadruple or increases 4 times.

1

2

3

4

5

6

7



Practice test 6 unit # 1



Correct



Unattempted



Incorrect



8/10

Q : The path of projectile in presence of air friction is more likely to be



Parabolic



Circle



Elliptical



Straight line

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT

4

5

6

7

8

9

10



Practice test 6 unit # 1



Correct



Unattempted



Incorrect



9/10

Q : In presence of air resistance, which one will be true for horizontal component of acceleration of projectile



$$a_x = 0$$



$$a_x = g$$



$$a_x \neq 0$$



$$a_x = -F_x$$

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT

4

5

6

7

8

9

10



Practice test 6 unit # 1



Correct



Unattempted



Incorrect



10/10

Q :

If a projectile is projected with initial speed 20 msec^{-1} at angle 75° with horizontal in presence of air friction, then which of the following may be the range of projectile



36 m



27 m



20 m



17 m

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT

4

5

6

7

8

9

10

www.saeedmdcat.com



FB) SAEED MDCAT

03418729745



QUIZZES

Class Test-1 (Unit # 1)



30 Questions



25 min

Topics

Velocity, Displacement-time graph,
Newton's laws of motion, Linear
Momentum, Law of conservation
of momentum, Projectile motion,
Characteristics of projectile motion, Time
of flight, Maximum height, Horizontal range,
Effect of Air Resistance in Projectile Motion

SAEED MDCAT

SAEED MDCAT TEAM

Start Quiz



SAEEDMDCAT



1/30



25 min



Hint

Q :

A train takes 1 hour to go from one station to the other. It travels at a speed of 30 kmh^{-1} for first half hour and at a speed of 50 kmh^{-1} for the next half hour. The average speed of the train is:



A 45 kmh^{-1}



B 35 kmh^{-1}



C 40 kmh^{-1}



D 30 kmh^{-1}

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



2/30



25 min



Hint

Q :

A boat is sent across a river with a velocity of 8kmh^{-1} . If the resultant velocity of boat is 10kmh^{-1} , then the velocity of the river is

A

12.8kmh^{-1}

B

6kmh^{-1}

C

8kmh^{-1}

D

10kmh^{-1}

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



3/30



25 min



Hint

Q :

Force of 5N acts on a body of 5kg for 5 sec then the rate of change in momentum is



25N



50N



5N



125N

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



4/30



25 min



Hint

Q : A cricket player catches a ball of mass 100 g and moving with a velocity of 25ms^{-1} . If the ball is caught 0.1s, the force of the ball exerted on the hand of the player is:



4N



40N



25N



250N

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



5/30



25 min



Hint

Q :

A cricket ball of mass 0.5 kg strikes a bat normally with a velocity of 30 ms^{-1} and rebounds with a velocity of 20 ms^{-1} in the opposite direction. The impulse of the force exerted by the ball on the bat is

A

0.5 Ns

B

25 Ns

C

1.0 Ns

D

50 Ns

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



6/30



25 min



Hint

Q : The rate of change of momentum of a body falling freely under gravity is equal to its

A

Impulse

B

Kinetic energy

C

Power

D

Weight

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



7/30



25 min



Hint

Q : A particle of mass m moving with velocity v strikes a stationary particle of mass $2m$ and sticks to it. The speed of the system will be?

A

$$\frac{v}{2}$$

B

$$2v$$

C

$$\frac{v}{3}$$

D

$$3v$$

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



8/30




25 min



Hint

Q : The displacement time graph for a moving particle is given below. The instantaneous velocity of the particle is negative at the point



A

D

B

F

C

C

D

E



SAEEDMDCAT



9/30



25 min



Hint

Q :

The displacement time graph of a particle moving with uniform velocity is

A

Parabola

B

Straight line

C

Circle

D

Hyperbola

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT

24 : 20



10/30



25 min



Hint

Q :

The displacement-time graphs of two particles A and B are straight lines making angles of 30° and 60° respectively with the time axis. If the velocity of A is v_A and that of B is v_B , the value of v_A/v_B is



1/2



$\frac{1}{\sqrt{3}}$



$\sqrt{3}$



1/3

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT

www.saeedmdcat.com

FB)SAEED MDCAT

03418729745

4

5

6

7

8

9

10



11/30




25 min




Hint

Q : To convert given displacement-time graph into distance time graph,




The corresponding distance time graph will be

A



B






C



03418729745

The corresponding distance time graph will be





12/30



25 min



Hint

Q : The slope of the tangent of a point on d-t graph gives the magnitude of

A

uniform velocity

B

Instantaneous velocity

C

average velocity

D

constant velocity

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



13/30



25 min



Hint

Q :

If d-t curve is a straight line, it shows that body is

A

average velocity

B

uniform velocity

C

both '(a)' and '(b)'

D

instantaneous velocity

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



14/30




25 min



Hint

Q :

A projectile is projected upwards with an angle of 45° with x-axis as shown in the figure such that its K.E at projection is 100J . What will be its K.E at the top?



0J



100J



25J



50J



SAEEDMDCAT



15/30



25 min



Hint

Q :

At which point for a projectile its kinetic energy is completely converted into potential energy

A

At point of projection

B

At the highest point

C

Point to hit the ground

D

Not possible

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



16/30



25 min



Hint

Q :

A 1000-kg airplane moves in straight flight at constant speed. The force of air friction is 1800 N. The net force on the plane is:



Zero



11800 N



1800 N



9800 N

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT

12

13

14

15

16

17

18



17/30



25 min



Hint

Q :

A projectile of mass m is thrown with a velocity v making an angle of 45° with the horizontal. The change in momentum from departure to arrival along vertical direction, is



2mv



$\sqrt{2}mv$



mv



$\frac{mv}{\sqrt{2}}$

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



18/30



25 min



Hint

Q : A ball is projected horizontally from the top of a cliff on the surface of the earth with a speed of 40ms^{-1} . Assuming that there is no air resistance, what will its speed be 3 s later?



A 30ms^{-1}



B 40ms^{-1}



C 50ms^{-1}



D 60ms^{-1}

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



19/30



25 min



Hint

Q :

A cannonball is fired and follows the parabolic path shown below. Air resistance is negligible. Point B is the highest point on the path and points A and C are at the same height



Which of the following best describes the direction of the velocity of the cannonball at point B?



to the right



down and to the right



up and to the right



up



20/30




25 min



Hint

A ball is fired and follows the parabolic path shown below. Air resistance is negligible. B is the highest point on the path and A and C are at the same height.



How do the accelerations of the ball at the points compare?

A

$$a_A < a_B < a_C$$

B

$$a_B < a_A < a_C$$

C

$$a_A = a_B = a_C$$

D

$$a_A = a_B < a_C$$



SAEEDMDCAT



20/30



25 min



Hint

Q :

cannonball is fired and follows the parabolic path shown below. Air resistance is negligible. Point B is the highest point on the path and points A and C are at the same height.



How do the accelerations of the ball at the three points compare?

A

$a_A < a_B < a_C$

B

$a_B < a_A < a_C$

C

$a_A = a_B = a_C$

D

$a_A = a_B < a_C$



21/30



25 min



Hint

Q : Two balls projected at 30° and 60° with same initial velocities. The ratio of their maximum heights is:



1:2



1:4



1:3



$1:\sqrt{2}$

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



22/30



25 min



Hint

Q : A projectile is projected with kinetic energy (K.E). If it has the maximum possible horizontal range, then its kinetic energy at the highest point will be:

A

K.E

B

0.75 K.E

C

0.5 K.E

D

0.25 K.E

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



23/30



25 min



Hint

Q : A body is thrown with a velocity of 9.8 m/s making an angle of 30° with the horizontal. It will hit the ground after a time



1 sec



2 sec



3 sec



1.5 sec

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



24/30



25 min



Hint

Q :

Time of flight of projectile is

A

$$\frac{v_i \sin \theta}{g}$$

B

$$\frac{v_i^2 \sin \theta}{g}$$

C

$$\frac{2v_i \sin^2 \theta}{g}$$

D

$$\frac{2v_i \sin \theta}{g}$$

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



25/30



25 min



Hint

Q : The maximum range of a projectile fired with some initial velocity is found to be 1000 m in the absence of wind and air resistance. The maximum height reached by the projectile is



250m



500 m



1000m



5000m

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



26/30



25 min



Hint

Q :

At which pair of angles the range of projectile would be same



30°, 60°



25°, 65°



15°, 75°



all of these

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



27/30



25 min



Hint

Q : Four projectiles are fired with the same velocity at angle, 25° , 40° , 55° and 70° with the horizontal. The range of projectile will be largest for the one projected at angle



70°



40°



55°



25°

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



28/30



25 min



Hint

Q :

Air resistance decreases the



horizontal component of velocity of a projectile



vertical component of velocity of a projectile



vertical component of acceleration of a projectile



none of these

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



29/30



25 min



Hint

Q :

When air resistance acts, acceleration during a fall will be

A

greater than g

B

less than g

C

equal to g

D

none of these

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



30/30



25 min



Hint

Q : The Factors affecting the flight path of a Projectile from the following are

A

Air Resistance

B

Gravity.

C

Angle of Release

D

All of these

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT



QUIZ RESULT

Class Test-1 (Unit # 1)



30



25 min



22-Feb-2021



0 sec



0/30



0.0%


SAEED MDCAT

Result Detail

SAEED MDCAT TEAM



SAEEDMDCAT





Class Test-1 (Unit # 1)

Q :

A train takes 1 hour to go from one station to the other. It travels at a speed of 30 kmh^{-1} for first half hour and at a speed of 50 kmh^{-1} for the next half hour. The average speed of the train is:

A

45 kmh^{-1}

B

35 kmh^{-1}

C

40 kmh^{-1}

D

30 kmh^{-1}

Explanation

As it is the equal time partition case so

$$V_{av} = \frac{v_1 + v_2}{2} = \frac{30 + 50}{2} = \frac{80}{2} = 40 \text{ kmh}^{-1}$$

1

2

3

4

5

6

7



Class Test-1 (Unit # 1)

Q :

A boat is sent across a river with a velocity of 8kmh^{-1} . If the resultant velocity of boat is 10kmh^{-1} , then the velocity of the river is

A

12.8kmh^{-1}

B

6kmh^{-1}

C

8kmh^{-1}

D

10kmh^{-1}

SAEED MDCAT

SAEED MDCAT TEAM

Explanation



SAEEDMDCAT

$$v_r = \sqrt{v_R^2 - v_B^2} = \sqrt{10^2 - 8^2} = 6 \text{ km}$$

1

2

3

4

5

6

7



Class Test-1 (Unit # 1)



Correct



Unattempted



Incorrect



3/30

Q :

Force of 5N acts on a body of 5kg for 5 sec then the rate of change in momentum is

A

25N

B

50N

C

5N

D

125N

Explanation

$$\frac{\Delta P}{\Delta t} = F$$
$$\frac{\Delta P}{\Delta t} = 5N$$

1

2

3

4

5

6

7



Class Test-1 (Unit # 1)



Correct



Unattempted



Incorrect



4/30

Q : A cricket player catches a ball of mass 100 g and moving with a velocity of 25ms^{-1} . If the ball is caught 0.1s, the force of the ball exerted on the hand of the player is:



4N



40N



25N



250N

Explanation

$$F = \frac{mv}{t} = \frac{(0.1)(25)}{0.1} = 25\text{N}$$



Class Test-1 (Unit # 1)



Correct



Unattempted



Incorrect



5/30

Q :

A cricket ball of mass 0.5 kg strikes a bat normally with a velocity of 30 ms⁻¹ and rebounds with a velocity of 20 ms⁻¹ in the opposite direction. The impulse of the force exerted by the ball on the bat is



0.5 Ns



25 Ns



1.0 Ns



50 Ns

Explanation

Impulse = change in momentum

$$I = mu - m(-v)$$

$$= m(u + v) = 0.5 \times (30 + 20) = 25 \text{ Ns}$$

1

2

3

4

5

6

7



Class Test-1 (Unit # 1)



Correct



Unattempted



Incorrect



6/30

Q : The rate of change of momentum of a body falling freely under gravity is equal to its



Impulse



Kinetic energy



Power



Weight

Explanation

$$F = \frac{\Delta p}{\Delta t} = w = mg$$

1

2

3

4

5

6

7



Class Test-1 (Unit # 1)



Correct



Unattempted



Incorrect



7/30

Q : A particle of mass m moving with velocity v strikes a stationary particle of mass $2m$ and sticks to it. The speed of the system will be?



$$\frac{v}{2}$$



$$2v$$



$$\frac{v}{3}$$



$$3v$$

Explanation



SAEEDMDCAT

$$mv + 0 = (m + 2m)v'$$
$$v' = \frac{mv}{3m} = \frac{v}{3}$$

1

2

3

4

5


6

7



Class Test-1 (Unit # 1)

Q : The displacement time graph for a moving particle is given below. The instantaneous velocity of the particle is negative at the point



A D

B F

C C

D E

Explanation



SAEEDMDCAT

At point E, the slope is decreasing negatively



Class Test-1 (Unit # 1)

The displacement time graph of a particle moving with uniform velocity is

- A Parabola
- B Straight line**
- C Circle
- D Hyperbola

Explanation

$\vec{d}-t$ graph represents that particle is moving with uniform velocity





Class Test-1 (Unit # 1)



Incorrect



10/30

Q :

The displacement-time graphs of two particles A and B are straight lines making angles of 30° and 60° respectively with the time axis. If the velocity of A is v_A and that of B is v_B , the value of v_A/v_B is

A

$1/2$

B

$\frac{1}{\sqrt{3}}$

C

$\sqrt{3}$

D

$1/3$

Explanation

SAEED MDCAT TEAM



SAEEDMDCAT

$$v_A = \tan 30^\circ \text{ and } v_B = \tan 60^\circ \therefore \frac{v_A}{v_B} = \frac{t}{t}$$



Class Test-1 (Unit # 1)



Incorrect



10/30

Q :

The displacement-time graphs of two particles A and B are straight lines making angles of 30° and 60° respectively with the time axis. If the velocity of A is v_A and that of B is v_B , the value of v_A/v_B is

A

$1/2$

B

$\frac{1}{\sqrt{3}}$

C

$\sqrt{3}$

D

$1/3$

Explanation

SAEED MDCAT TEAM




SAEEDMDCAT

$$= \tan 60^\circ \therefore \frac{v_A}{v_B} = \frac{\tan 30^\circ}{\tan 60^\circ} = \frac{1/\sqrt{3}}{\sqrt{3}} = \frac{1}{3}$$




Class Test-1 (Unit # 1)

Q : To convert given displacement-time graph into distance time graph,




The corresponding distance time graph will be

A



B



C





Class Test-1 (Unit # 1)



Correct



Unattempted



Incorrect



12/30

Q : The slope of the tangent of a point on d-t graph gives the magnitude of



uniform velocity



Instantaneous velocity



average velocity



constant velocity

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT

9

10

11

12

13

14

15



Class Test-1 (Unit # 1)



Correct



Unattempted



Incorrect



13/30

Q :

If d-t curve is a straight line, it shows that body is

A

average velocity

B

uniform velocity

C

both '(a)' and '(b)'

D

instantaneous velocity

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT

9

10

11

12


13

14

15



Class Test-1 (Unit # 1)



A

0J

B

100J

C

25J

D

50J

Explanation

$$\begin{aligned} K.E &= K.E_{\max} \cos^2 \theta \\ &= K.E_{\max} \left(\frac{1}{\sqrt{2}} \right)^2 \\ &= 100 \times \frac{1}{2} \\ &= 50J \end{aligned}$$



Class Test-1 (Unit # 1)



Incorrect



15/30

Q :

At which point for a projectile its kinetic energy is completely converted into potential energy

A

At point of projection

B

At the highest point

C

Point to hit the ground

D

Not possible

Explanation

There is no point in the path of a projectile where K.E is completely converted to P.E every point have some amount of K.E and P.E.

9

10

11

12

13

14

15



Class Test-1 (Unit # 1)

Q :

A 1000-kg airplane moves in straight flight at constant speed. The force of air friction is 1800 N. The net force on the plane is:

A

Zero

B

11800 N

C

1800 N

D

9800 N

Explanation

SAEED MDCAT TEAM

$$v = \text{constant} \Rightarrow \Delta v = 0$$

$$a_{\text{net}} = 0 \Rightarrow F_{\text{net}} = 0$$

$$v = \text{constant} \Rightarrow \Delta v = 0$$

$$a_{\text{net}} = 0 \Rightarrow F_{\text{net}} = 0$$



Class Test-1 (Unit # 1)

horizontal. The change in momentum from departure to arrival along vertical direction, is

A

$$2mv$$

B

$$\sqrt{2}mv$$

C

$$mv$$

D

$$\frac{mv}{\sqrt{2}}$$

Explanation

Change in momentum is the product of force and time

$$\Delta p = mg \times \frac{2v \sin \theta}{g}$$

$$= 2mv \sin \theta$$

$$= 2mv \sin 45^\circ = \frac{2mv}{\sqrt{2}} = \sqrt{2}mv$$



Class Test-1 (Unit # 1)



Correct



Unattempted



Incorrect



18/30

Q : A ball is projected horizontally from the top of a cliff on the surface of the earth with a speed of 40ms^{-1} . Assuming that there is no air resistance, what will its speed be 3 s later?

A

30ms^{-1}

B

40ms^{-1}

C

50ms^{-1}

D

60ms^{-1}


Explanation

$$v = \sqrt{v_x^2 + g^2 t^2} = \sqrt{(40)^2 + (10)^2 (3)^2}$$
$$v = \sqrt{1600 + 900} = \sqrt{2500} = 50\text{ms}^{-1}$$



Class Test-1 (Unit # 1)

points A and C are at the same height



Which of the following best describes the direction of the velocity of the cannonball at point B?

A

to the right

B

down and to the right

C

up and to the right

D

up


Explanation

at highest point only horizontal velocity acts which having direction horizontally towards right according to given diagram



Class Test-1 (Unit # 1)

cannonball is fired and follows the parabolic path shown below. Air resistance is negligible. Point B is the highest point on the path and points A and C are at the same height.



How do the accelerations of the ball at the three points compare?

A

$$a_A < a_B < a_C$$

B

$$a_B < a_A < a_C$$

C

$$a_A = a_B = a_C$$

D

$$a_A = a_B < a_C$$

Explanation

Acceleration in projectile motion at every point is equal to g which has direction directed towards downward.



Class Test-1 (Unit # 1)



Correct



Unattempted



Incorrect



21/30

Q : Two balls projected at 30° and 60° with same initial velocities. The ratio of their maximum heights is:



1:2



1:4



1:3



$1:\sqrt{2}$

Explanation

As

$$H_{max} = \frac{v_i^2 \sin^2 \theta}{2g}, \text{ As } v_i \text{ is same so } \frac{H_1}{H_2} = \frac{\sin^2 \theta_1}{\sin^2 \theta_2}$$



Class Test-1 (Unit # 1)



Correct



Unattempted



Incorrect



21/30

Q : Two balls projected at 30° and 60° with same initial velocities. The ratio of their maximum heights is:



1:2



1:4



1:3



$1:\sqrt{2}$

Explanation

As v_i is same so $\frac{H_1}{H_2} = \frac{\sin^2 \theta_1}{\sin^2 \theta_2} = \frac{(\sin 30)^\circ}{(\sin 60)^\circ} = 1:3$



Class Test-1 (Unit # 1)



Correct



Unattempted



Incorrect



22/30

Q : A projectile is projected with kinetic energy (K.E). If it has the maximum possible horizontal range, then its kinetic energy at the highest point will be:



K.E



0.75 K.E



0.5 K.E



0.25 K.E

Explanation

$$K.E_H = K.E_1 \cos^2 \theta$$

$$K.E_H = K.E_1 \cos^2 45^\circ = 0.5 K.eE$$



Class Test-1 (Unit # 1)



Correct



Unattempted



Incorrect



23/30

Q : A body is thrown with a velocity of 9.8 m/s making an angle of 30° with the horizontal. It will hit the ground after a time



1 sec



2 sec



3 sec



1.5 sec

Explanation

$$t = \frac{2v_i^2 \sin \theta}{g} = \frac{2(9.8) \sin 30}{(9.8)} = 1 \text{ sec}$$



Class Test-1 (Unit # 1)



Incorrect



24/30

Q :

Time of flight of projectile is

A

$$\frac{v_i \sin \theta}{g}$$

B

$$\frac{v_i^2 \sin \theta}{g}$$

C

$$\frac{2v_i \sin^2 \theta}{g}$$

D

$$\frac{2v_i \sin \theta}{g}$$

SAEED MDCAT TEAM

Explanation



SAEEDMDCAT

$$\frac{2v_i \sin \theta}{g}$$

is the time of flight of projectile.



Class Test-1 (Unit # 1)



Correct



Unattempted



Incorrect



25/30

Q : The maximum range of a projectile fired with some initial velocity is found to be 1000 m in the absence of wind and air resistance. The maximum height reached by the projectile is



250m



500 m



1000m



5000m

Explanation

$$R \tan \theta = 4H$$

$$\theta = 45^\circ$$

$$H = \frac{R}{4} = \frac{1000}{4} = 250 \text{ m}$$

24

25

26

27

28

29

30



Class Test-1 (Unit # 1)



Correct



Unattempted



Incorrect



26/30

Q :

At which pair of angles the range of projectile would be same



30°, 60°



25°, 65°



15°, 75°



all of these

Explanation



$$\theta_1 + \theta_2 = 90^\circ$$

$$R\theta_1 = R\theta_2$$

SAEEDMDCAT



Class Test-1 (Unit # 1)



Correct



Unattempted



Incorrect



27/30

Q : Four projectiles are fired with the same velocity at angle, 25° , 40° , 55° and 70° with the horizontal. The range of projectile will be largest for the one projected at angle



70°



40°



55°



25°

SAEED MDCAT

SAEED MDCAT TEAM

Explanation



SAEEDMDCAT

Range will be maximum at that angle which is nearest to 45°

24

25

26

27

28

29

30



Class Test-1 (Unit # 1)



Correct



Unattempted



Incorrect



28/30

Q :
Air resistance decreases the



horizontal component of velocity of
a projectile



vertical component of velocity of
a projectile



vertical component of acceleration of
a projectile



none of these

SAEED MDCAT

SAEED MDCAT TEAM



SAEEDMDCAT

24

25

26

27

28

29

30



Class Test-1 (Unit # 1)



Incorrect



29/30

Q :

When air resistance acts, acceleration during a fall will be

A

greater than g

B

less than g

C

equal to g

D

none of these

Explanation

acceleration during a fall will be less than g because air resistance affects the motion of the falling objects by slowing it down

24

25

26

27

28

29

30



Class Test-1 (Unit # 1)



Correct



Unattempted



Incorrect



30/30

Q : The Factors affecting the flight path of a Projectile from the following are



Air Resistance



Gravity.



Angle of Release



All of these

Explanation

all factors given are responsible for the flight of projectile

24

25

26

27

28

29

30

SAEED MDCAT

TRUE HERO ALWAYS WINS

National MDCAT 2020 Results

190+

35 Students

185+

218 Students

180+

677 Students



SAEED MDCAT
SAEED MDCAT TEAM
f SAEEDMDCAT

SAEED MDCAT
JOIN US FOR FREE
TO SECURE YOUR FUTURE

Designed by: Usman Arts



SAEEDMDCAT